Version of Record: https://www.sciencedirect.com/science/article/pii/S0079661116000057
Manuscript_61f7b14656d59e5f2a27b3f0854bbc87

A hierarchical approach to defining marine heatwaves

Alistair J. Hobday[1]
Lisa V. Alexander[2,3]
Sarah E. Perkins[2,3]
Dan A. Smale[4,5]
Sandra C. Straub[5]
Eric C. J. Oliver[2,6]
Jessica Benthuysen[7]
Michael T. Burrows[8]
Markus G. Donat[2,3]
Ming Feng[9]
Neil J. Holbrook[2,6]
Pippa J. Moore[10]
Hillary A. Scannell[11,12]
Alex Sen Gupta[3]
Thomas Wernberg[5]

1. CSIRO Oceans and Atmosphere, Hobart, Tasmania, 7000, Australia
2. ARC Centre of Excellence for Climate System Science, The University of New South Wales, Sydney, Australia
3. Climate Change Research Centre, The University of New South Wales, Sydney, Australia
4. Marine Biological Association of the United Kingdom, The Laboratory, Citadel Hill, Plymouth PL1 2PB, UK
5. UWA Oceans Institute and School of Plant Biology, The University of Western Australia, Crawley 6009 Western Australia, Australia
6. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia
7. Australian Institute of Marine Science, Townsville, Queensland, Australia
8. Department of Ecology, Scottish Association for Marine Science, Scottish Marine Institute, Oban, Argyll, PA37 1QA, Scotland, UK.
9. CSIRO Oceans and Atmosphere, Perth, Western Australia, Australia
10. Institute of Biological, Environmental and Rural Sciences, Aberystwyth University, Aberystwyth SY23 3DA, UK
11. School of Oceanography, University of Washington, Seattle, WA, USA
12. NOAA/Pacific Marine Environmental Laboratory, Seattle, WA, USA

Version: December 30, 2015

Progress in Oceanography

© 2016 published by Elsevier. This manuscript is made available under the Elsevier user license
https://www.elsevier.com/open-access/userlicense/1.0/

Abstract

Marine heatwaves (MHWs) have been observed around the world and are expected to increase in intensity and frequency under anthropogenic climate change. A variety of impacts have been associated with these anomalous events, including shifts in species ranges, local extinctions and economic impacts on seafood industries through declines in important fishery species and impacts on aquaculture. Extreme temperatures are increasingly seen as important influences on biological systems, yet a consistent definition of MHWs does not exist. A clear definition will facilitate retrospective comparisons between MHWs, enabling the synthesis and a mechanistic understanding of the role of MHWs in marine ecosystems. Building on research into atmospheric heatwaves, we propose both a general and specific definition for MHWs, based on a hierarchy of metrics that allow for different data sets to be used in identifying MHWs. We generally define a MHW as a prolonged discrete anomalously warm water event that can be described by its duration, intensity, rate of evolution, and spatial extent. Specifically, we consider an anomalously warm event to be a MHW if it lasts for five or more days, with temperatures warmer than the 90th percentile based on a 30-year historical baseline period. This structure provides flexibility with regard to the description of MHWs and transparency in communicating MHWs to a general audience. The use of these metrics is illustrated for three 21st century MHWs; the northern Mediterranean event in 2003, the Western Australia 'Ningaloo Niño' in 2011, and the northwest Atlantic event in 2012. We recommend a specific quantitative definition for MHWs to facilitate global comparisons and to advance our understanding of these phenomena.

Keywords: extreme events; sea surface temperature; anomalous events; temperature anomaly; heatwaves

1. Introduction - Marine heatwaves and their ecological impact

Ecosystems around the world have responded to anthropogenic climate change, with major implications for ecological goods and services (Rosenzweig et al. 2008). Links between a changing climate, shifts in species distributions, and the structure of communities and ecosystems have been documented convincingly for many taxa across many regions (Parmesan and Yohe 2003; Rosenzweig et al. 2008; Poloczanska et al. 2013). Concurrent with these observations, predictions of how species distribution and biodiversity will respond to continued climate change have been developed (e.g. Cheung et al. 2009; Engler et al. 2011; Sen Gupta et al. 2015). However, in conjunction with a distinct long-term warming signal (an increase in mean temperature at a location), the frequency and intensity of extreme temperature events are also increasing (Perkins et al. 2012) as a consequence of anthropogenic climate change (IPCC 2012; Coumou and Ramstorf 2012). It is clear that discrete climatic events can drive step-wise changes in species distributions and, ultimately, ecosystem structure and functioning (Wernberg et al. 2013). Storms, droughts, floods and heatwaves - prolonged period where temperatures are substantially hotter than normal - can have catastrophic effects on terrestrial ecosystems (Jentsch et al. 2007; Smith 2011), with significant socio-economic ramifications. As such, understanding and predicting biological responses to short-term extreme events, rather than long-term change, is becoming increasingly important, although event-based research still lags behind trend-based work (Jentsch et al. 2007).

Extreme climatic events are important in determining ecosystem structure (Jentsch et al. 2007), however, the majority of our current understanding stems from the study of terrestrial ecosystems. Investigation of marine ecosystems is important, as they play a central role culturally, socially and economically in the lives of most people (Richardson and Poloczanska 2008; Bennett et al. 2015). Marine ecosystems, like their terrestrial counterparts, are strongly influenced by extreme climatic events, including heatwaves (Garrabou et al. 2009; Wernberg et al. 2013), cold snaps (Firth et al. 2011), storms (De'ath et al. 2012) and floods (Gillanders and Kingsford 2002), which are driven by complex physical processes interconnected in the climate system and interacting across a hierarchy of spatial and temporal scales (Trenberth 2012; Feng et al. 2014).

Marine heatwaves (MHWs), which can be caused by a combination of atmospheric and oceanographic processes, have a strong influence on marine ecosystem structure and function. For example, in the boreal summer of 2003 an atmospheric heatwave over northwestern Europe led to enhanced rates of air-sea heat flux into the northern Mediterranean Sea, which in combination with weak winds led to regional-scale thermal stratification and warming anomalies of 2-3°C in surface waters (Garrabou et al. 2009). This MHW had profound ecological impacts that included widespread mortality of benthic invertebrates (Garrabou et al. 2009) and loss of seagrass meadows (Marba and Duarte 2010). More recently, during the austral summer of 2011, a MHW off Western Australia (a so-called 'Ningaloo Niño') was largely driven by atmospheric and oceanographic processes associated with the strong 2010/11 La Niña, which led to anomalous advection of warm tropical waters poleward into temperate regions (Feng et al. 2013; Benthuysen et al. 2014). This Western Australia MHW caused major shifts in benthic ecosystem structure and functioning in a tropical-temperate transition zone, through widespread mortality of cool-water habitat forming species (Wernberg et al. 2013; Smale and Wernberg 2013), and impacted a valuable fishery (Caputi et al. 2015). During a 2012 MHW in the northwest Atlantic, rapid shifts in geographical distributions and phenology were observed for several marine species, including those targeted by regional fisheries (Mills et al. 2013). These ecological responses led to altered fishing practices (longfin squid) and harvest patterns (lobsters), with significant political and economic ramifications (Mills et al. 2013).

It is clear that MHWs, which may increase in frequency and magnitude as a result of anthropogenic climate change (IPCC 2012), are important events that can cause rapid changes in biodiversity patterns and ecosystem structure and functioning. Apart from the physical drivers of short-term temperature variability and extremes, there is a pressing need to examine the characteristics of MHWs, and their biological impacts, within a coherent and comparable framework.

1.1. Defining extreme temperatures in marine systems

Previous ecological studies have used metrics to assess extreme thermal stress in the marine environment (**Table 1**). For example, Sorte et al. (2010) adopted the definition of Meehl and Tebaldi (2004) in which marine heatwaves were defined as a period of at least three to five days during which mean or maximum temperature anomalies were at least 3 – 5°C above normal, while Selig et al. (2010) used thermal stress anomalies (TSAs – see **Table 1**).

Recently, Marba et al. (2015) used SST percentile thresholds for a Mediterranean-focused meta-analysis of MHW impacts, however MHWs are also often described using vague definitions (e.g. statements such as "warmer than average") and most temperature anomalies are generated from monthly datasets, thus smoothing out shorter but generally more intense events. The majority of marine extreme climate event metrics have been developed to monitor and predict coral bleaching, which is the most advanced field of thermal stress-related marine ecology (Donner et al. 2005; Spillman and Alves 2009). Such metrics generally include the effect of extreme event duration and magnitude of temperature anomalies. Beyond coral reef research there is limited consistency regarding how MHW metrics are applied or how useful they are in ecological applications.

1.2. Parallels with atmospheric heatwave definitions

Global initiatives over the last decade have sought to define standard metrics for atmospheric heatwaves and extreme temperatures, primarily under the auspices of the Expert Team on Climate Change Detection and Indices (ETCCDI[1]; Zhang et al. 2011), to allow comparative analyses across regions. The general definition of atmospheric heatwaves is a prolonged period where temperatures are substantially hotter than normal (Perkins and Alexander 2013). Observations of atmospheric extreme events have had considerably more attention over the last decade compared with marine events (e.g. Meehl and Tebaldi 2004; Fischer and Shär 2010; Schoetter et al. 2014), but the absence of a pre-defined framework has seen atmospheric events defined by a plethora of metrics, most of which are unique to a particular purpose or study. The existing metrics are generally simplistic, accounting for only anomaly, magnitude, duration or frequency (Frich et al. 2002; Alexander et al. 2006; Perkins 2011). A few studies, however, have attempted to develop more complex metrics that take into account multiple factors (Della-Marta et al. 2007; Vautard et al. 2013; Russo et al. 2014). In parallel to the climate research community, impact-focused research groups (such as the health sector) have defined an additional suite of heatwave metrics. While these indices also measure the severity of heatwaves, they are configured to relate to thresholds that are specific to a particular application (e.g. metrics for human health purposes: Fanger 1970; Steadman 1984; Mayer and Hoppe 1987). Such metrics often require more than just basic temperature data, making it difficult to derive most impact metrics from regional climatological data. Moreover, the specific nature of each metric to a particular impact reduces its applicability to

[1] A joint initiative of the World Meteorological Organisation Commission for Climatology/World Climate Research Programme/ and the Joint WMO-IOC Technical Commission for Oceanography and Marine Meteorology (JCOMM).

another sector, even with similar purposes (Perkins and Alexander 2013). This wide range of metrics within and across these communities also means that different data are required to apply different atmospheric heatwave definitions, which inhibits consistent measurements both spatially and temporally. The lack of consistency in data availability and atmospheric heatwave calculations has made a general assessment of the drivers of these events and their impacts extremely challenging. These limitations have resulted in an assessment for observed trends in atmospheric heatwaves of only *medium confidence* in the IPCC Special Report on Extremes (IPCC 2012) and the IPCC Fifth Assessment Report (IPCC 2013).

In an attempt to overcome these issues, Perkins and Alexander (2013) presented a working framework to define atmospheric heatwaves and address the issues of inconsistency and assigning confidence. The framework considers what metrics can be derived with statistical rigor from meteorological data, and what characteristics are important for a range of impacts sectors. Based on these criteria, Perkins and Alexander (2013) define an atmospheric heatwave when at least three consecutive days exceed a calendar day threshold defined as the 90^{th} percentile value for temperature. Using a 'day-specific' threshold allows for the detection and measurement of events at all times of the year (i.e. a heatwave can occur in winter with a lower absolute value than might occur in summer), and a percentile-based threshold allows for the measurement of heatwaves across locations that differ in variability. An event is characterized in terms of its duration and intensity, and summary statistics such as the number of discrete events, sum of heatwave days and peak intensity can be calculated for a season or period of interest. The success of the framework is evident in understanding changes in global observed atmospheric heatwaves (Perkins et al. 2012) and future projections from numerical climate models (Cowan et al. 2014). It also supported a finding of *high confidence* in observed increasing trends in heatwave frequency in Europe, Asia and Australia (IPCC 2013).

While the framework constructed by Perkins and Alexander (2013) has achieved a consistent approach to characterising atmospheric heatwaves, the study of atmospheric heatwaves would have likely been more successful if common definitions had been derived earlier in the study of atmospheric heatwaves. This success would have been further heightened by incorporating levels of metric flexibility and ease of communication. Therefore, there is great potential for the marine community to apply the lessons learned from the atmospheric community in the definition of MHWs.

2. A hierarchical definition of marine heatwaves

From the lessons learned in atmospheric studies, and following Perkins and Alexander (2013), we propose a definition for MHWs that can be used for comparative studies with regional and biological applications. Minor differences to the atmospheric definition (minimum duration and minimum time between events) were implemented because of the naturally longer time scales of ocean variability with regard to atmospheric variability, as explained below. Qualitatively, we propose the definition of a MHW as **a discrete prolonged anomalously warm water event** in a particular location. From examples such as the 2003 MHW in the northern Mediterranean Sea (Garrabou et al. 2009), the 2011 Ningaloo Niño in Western Australia (Feng et al. 2013) and the 2012 MHW in the northwest Atlantic (Mills et al. 2013; Chen et al. 2014), it is clear that the atmosphere, land surface and ocean all may have important driving roles in how and where these prolonged heat events play out. However the relative importance of these drivers varies amongst events. Therefore, the qualitative definition does not assume any particular heatwave driver nor does it assume that the MHW has any specific impact. However, it does provide a flexible definition that can be specifically targeted towards end-user applications such as coral reef monitoring or fisheries management. In these situations, identification and quantification of heatwave events provides an opportunity to understand and manage impacts, such as when the 2011 Ningaloo Niño decimated commercially important crustacean and mollusc stocks in Western Australia (Pearce et al. 2011; Hodgkinson et al. 2014).

The qualitative definition of a MHW applies to ocean regions (including subsurface waters, estuarine, or enclosed seas such as the Mediterranean Sea or Baltic Sea), but may have limited applications in intertidal zones, where ecological responses to high sea temperatures are mediated by air temperature, precipitation and atmospheric conditions (Helmuth et al. 2006). Under this definition, a MHW can be caused by a combination of atmospheric forcing (e.g. heating) and oceanic conditions (e.g. faster advection or advection of unusually warm water). The MHW should be defined relative to a baseline period (climatology) and a particular time of the year from which the intensity, duration and spatial extent of the MHW could be defined. This also means that a MHW is not just limited to the warmer months, since for some biological applications the consideration of heatwaves in colder months is essential. For example, the reproductive cycle of several seaweed species involves reproduction in colder seasons, and during these seasons the propagules and early post-

settlement stages are in general more susceptible to thermal stress than adults (Santelices 1990; Lotze et al. 2001). For the fucoid *Scytothalia dorycarpa*, Andrews et al. (2014) showed that post-settlement juvenile survivorship strongly depended on temperature, with highest survivorship in the coldest treatment, and elevated or complete mortality more likely under elevated temperature. In this case, a MHW in a cold season could lead to suppressed or failed recruitment of habitat-forming seaweeds.

While this qualitative definition provides flexibility in the way in which a MHW can be defined across multiple end users for their particular application, it does not allow for empirical comparisons of the characteristics of MHWs across different events in space and time. For intercomparisons, the general qualitative definition of 'anomalously warm', 'discrete', and 'prolonged' can be quantified:

- '**anomalously warm**': A MHW must be defined relative to a baseline climatology (see recommendation section). Based on other studies of ocean drivers (e.g. El Niño-Southern Oscillation), which have long time scales of variability, a period of 30 years[2] is recommended to define a baseline temperature climatology, wherever possible. This is almost the full period of recorded satellite sea surface temperature observations. The climatology will be defined relative to the time of year, using all data within an 11-day window centred on the time of year from which the climatological mean and threshold are calculated. Limitations, in terms of length, quality, consistency, resolution and availability may restrict this method for some applications. For studies using remotely sensed data, where availability begins in the 1980s and 1990s for sea surface temperature and sea surface height, respectively, the climatological period might have to be shorter, and users should explicitly define their period accordingly. A MHW should be defined relative to a high percentile threshold (e.g. 90%). A percentile threshold is recommended rather than an absolute value above the climatological value as the magnitude of variability across a range of timescales varies considerably by region. An absolute threshold (e.g. 2°C anomaly) would only be relevant in terms of impacts in some regions but not in others (e.g. due to species acclimation). Moreover, by using a percentile rather than standard deviation definition no assumption is made regarding the underlying distribution of anomalies. Users should also be cognisant of biases that might be

[2] Guide to Climatological Practices, WMO-No. 100

introduced at the start and end of the base period when calculating threshold exceedances, and in such cases a bootstrapping procedure such as that defined by Zhang et al. (2005) might be employed to calculate percentiles from subsets of the data when a long time series is available.

- '**prolonged'**: In the marine environment, the definition should be relevant to ecological processes and thresholds (based on evidence of impact), but for each process this threshold may be different. Our general recommendation is that the MHW needs to persist for at least five days. A sensitivity analysis was performed using high-resolution (1/4°), global, daily SSTs from the Advanced Very High Resolution Radiometer (AVHRR) satellite data (NOAA OI SST V2; Reynolds et al. 2007) and it was found that for durations shorter than five days there were many more MHWs in the tropical regions than elsewhere, while for durations longer than five days there were often many regions with fewer than one MHW per year, on average. Therefore, we recommend five days as a balance to achieve relatively uniform global MHW counts under current climatic conditions.
- '**discrete**': A MHW event is discrete with well-defined start and end times. However, in our proposed definition and in common with atmospheric heatwaves, gaps between events of two days or less with subsequent five day or more events will be considered as a continuous event. For example, five anomalously warm days followed by two cool and then six anomalously warm days would be defined as an 13 day MHW event [5_{hot},2_{cool},6_{hot}]. In contrast, five anomalously warm days, followed by one cool day, and then two more anomalously warm days would be defined as a five day event [5_{hot},1_{cool},2_{hot} = 5 MHW days]; as would the converse [2_{hot},1_{cool},5_{hot}]. A sequence of five anomalously warm days followed by four cool days and then six anomalously warm days [5_{hot},4_{cool},6_{hot}] would be defined as two MHW events, one of five days duration, and one of six days duration.

2.1. Measurement of marine heatwaves

MHWs can be identified at any point in the ocean based on quantitative refinement of the qualitative definition provided earlier. For intercomparisons, a standard MHW definition, calculated in exactly the same way and using the same metrics and processing methods, is required. We suggest that the previous values be used as a starting point, but could be modified for a particular region or purpose. A set of summary statistics can be derived for each MHW including, for example, its intensity, duration, frequency and spatial extent. We

propose that a hierarchical set of such metrics be used to uniquely describe MHWs (**Figure 1; Table 2**). A hierarchy is useful as different temperature datasets, based on their spatial and temporal resolution, have different abilities to provide different metrics. Primary metrics allow for the most general comparison between duration and magnitude (intensity). For example, for a MHW, duration is defined as the period over which the temperature is greater than the seasonally varying threshold value (also defined in **Table 2**), while cumulative intensity (i_{cum}) is the integral of intensity over the duration of the event, and is equivalent to previously used metrics such as DHDs. Secondary metrics distinguish the temporal trend (i.e. the rate of event onset and decline) and spatial extent of the MHW. Tertiary metrics are very specific to the system under investigation, and include preconditioning environmental conditions, although we do not formally define these conditions. This hierarchy allows some flexibility in the reporting of MHWs, particularly for non-scientific audiences. Measures such as duration and intensity are easily understood, while rates of onset and decline and cumulative intensity may require additional explanation. This set of metrics allows different MHW events to be uniquely described and compared (**Figure 2**). The MHW definition as used in this manuscript has been implemented as a free software package in Python that calculates all the metrics for a provided time series (marineHeatWaves, http://github.com/ecjoliver/marineHeatWaves).

As described earlier, a number of MHWs have been recorded over recent decades but have been mainly described in general terms as abnormally warm or several degrees above the mean. Comparison of these events across marine environments would be possible by calculating one or more common metrics to all past MHWs. This, in turn, would allow a characterisation based on the hierarchical classifications of metrics, placing the events in a historical context. As an example, three better-known MHW regions are compared here to illustrate the use of these metrics (**Figure 3**). The metrics for each location were derived from NOAA OI SST, using code implemented in Python (available from https://github.com/ecjoliver/marineHeatWaves). Each region has numerous MHW events that meet our criteria based on the duration and intensity of each event. For example, the location examined off Western Australia (**Figure 3a)** has experienced 59 MHW events (duration of five days or more) between 1982 and 2014, with the longest MHW lasting for 95 days (13 May 1999 – 15 August 1999) with a maximum (i_{max}), mean (i_{mean}) and cumulative intensity (i_{cum}) of 3.60°C, 2.50 °C and 237°C days above the climatological mean, respectively. By way of comparison, the 2011 event was the largest event according to i_{max} (6.50°C) and i_{mean}

(3.21°C), and the second largest MHW after the 1999 event according to duration (60 days) and i_{cum} (192°C days).

For the Mediterranean Sea location, a total of 70 events were identified (**Figure 3b)**. Different MHW events had the longest duration (2014), highest maximum intensity (2008), and highest mean intensity (2003). The 2003 MHW was the largest event based on the i_{mean} (4.06°C; not shown in Figure 3b) and lasted 30 days by our definition (2 June – 1 July) with an i_{max} of 5.02°C. The most intense event was in 2008 (i_{max} = 5.05) lasting only 9 days (26 June to 4 July) with an i_{mean} of 3.87°C. The longest event was still ongoing at the time of analysis, with 110 days (13 September - 31 December 2014 – the end of the dataset), with a lower i_{max} (3.31°C) and i_{mean} (2.51°C), but the highest i_{cum} (276°C days).

For the selected northwest Atlantic location, 67 events were identified (**Figure 3c)**, with the longest MHW of duration 187 days (31 July 2012 - 2 February 2013) with an i_{max} and i_{mean} of 4.00°C and 2.37°C respectively. The i_{cum} for this MHW was 443°C days, the highest for any at this location. An earlier event, lasting 21 days (4-24 July 2010), had the highest mean intensity (3.05°C) in the period considered, but a lower maximum intensity (i_{max}=4.24°C) than a 56 day event in 2012 (10 April – 4 June) where the i_{max} was 4.89°C. This latter event is the 2012 northwest Atlantic event discussed in the literature (Mills et al. 2013; Chen et al. 2014). The corresponding i_{mean} and i_{cum} for this 56 day event was 2.59°C and 145 °C days, respectively. Note that the 56 and 187 day events in 2012 are considered distinct, as the temperature dropped below the threshold for at least 3 days (5 days) between the two events.

The collective analysis of the three case study regions demonstrates the need for a diversity of metrics to describe different MHW features. As each of the MHWs is defined by a set of metrics (**Table 2)**, approaches such as principal component analysis can be undertaken to characterise and identify types of MHWs. The metrics for each location may also be used to examine how the frequency of events has changed over time by analysing individual events (e.g. **Figure 2**), or the total number of MHW days in each year. Finally, the spatial extent of MHWs can be calculated from gridded datasets (e.g. NOAA OI SST) with the analysis of temperature time series repeated for each point in a spatial grid in the wider region of interest. The area where the threshold is exceeded is summed for each day to provide a daily MHW area for each day. These metrics could in turn be used to explore the impact of MHWs on regional biology. As evident from published studies outlined above, persistent and intense

MHWs have led to widespread and notable ecological impacts, analogous to atmospheric heatwaves. With these consistent set of metrics, comparative analyses, including linking ecological impacts to specific MWH characteristics, can be undertaken. While we have included 'preconditioning' as a tertiary metric in our hierarchy (**Table 2**), we do not expand further here, as these metrics will likely be specific to particular habitats, regions and species via potential local adaptation to extremes (Palumbi et al. 2014). When researchers describe MHWs in the future, consideration of preconditioning, such as a period of warm, but not anomalous, conditions may provide additional insight into ecological or human impacts of the focal MHW. Their inclusion in our hierarchy thus represents a placeholder to be informed by more studies on preconditioning and may be expanded or discarded in the future.

2.2. Datasets matter in defining heatwaves

While a consistent framework to measure MHWs is important, end-users need to be aware that different datasets may provide substantially different heatwave information despite the use of the same metrics. This is generally due to the resolution of the data, but can also relate to other issues of quality, consistency and instrumentation. Datasets with a high spatial and temporal resolution have more variability than those aggregating across larger areas or based on (smoother) longer time means (Smale and Wernberg 2009). An example of the variation that arises from using different datasets for MHW identification is given in **Figure 4**, which shows the development of the Ningaloo Niño in 2011 from the reconstructed monthly and weekly Reynolds SST dataset (Reynolds et al. 2002), a daily satellite product (NOAA OISST product; Reynolds et al. 2007), and an hourly *in situ* data logger. All datasets have a similar profile of the evolution of summer and the MHW including the rate of onset and decline, the duration of the event (measured in months), and a warm period preceding the main heatwave. However, the variability in SST magnitude clearly differs between the four datasets, and would result in different metrics of heatwave intensity. The reconstructed SST products have the smallest variability, due to the coarse spatial (1° degree grid) and temporal (monthly and weekly) resolution. This is followed by the daily satellite data, which are finer in spatial (0.25° degree grid) and temporal (daily) resolution. The high temporal resolution provided by the *in situ* logger data reveals higher temporal SST variability, but a lower peak intensity than the daily satellite dataset, consistent with previous analysis of sub-surface *in situ* and daily satellite data in this region (Smale and Wernberg 2009). It is clear that weekly variability in the logger data is smoothed at monthly scales, thus decreasing intensity by including non-

heatwave days and weeks. Lower spatial resolution data results in reduced intensity because neighbouring non-heatwave areas are included in mean values.

Not only can different datasets generate different values for the same metric, certain indices may simply not be appropriate or derivable from some data sources. **Table 3** gives an indication of when the indices outlined in the framework may or may not be applicable, and the quality they would provide. For example, *in situ* data (such as the logger data described above) can provide high frequency information for the more accurate calculation of intensity and duration (if measured in days). However, these local data would not provide an estimate of the spatial extent of a MHW. In contrast, gridded products, such as satellite-derived SSTs and reconstructed daily datasets, allow greater spatial inferences. Model data, in the forms of global and regional models, and reanalysis products, if at a daily scale, may be used for the calculation of all MHW metrics (**Table 3**). However, resolution may be reduced due to the coarseness of spatial grids and, in the case of regional models, the domain they cover. While useful for other purposes, paleo proxies and traditional ecological data can, at best, only provide quantitative information on long-lived MHWs (e.g. Zinke et al. 2014). A number of other considerations listed in **Table 3**, including record length, temporal resolution, whether the data have been quality-controlled, and spatio/temporal consistency, should help end-users evaluate what metrics can be derived from a particular product. Such considerations and measurement qualities are indicative only and should be applied to a dataset each time it is used for the measurement of MHWs.

Many of the MHW metrics can be calculated from gridded products, such as SST datasets, reconstructed observational data, and model/reanalysis data. These provide generally similar quality metrics (**Table 3**). We recommend that the highest quality data available should be used when calculating MHWs and where possible compared to *in situ* data (also of high-quality, e.g. Smale and Wernberg 2009) (e.g. **Figure 4**). While coarser resolution datasets may provide information about larger areas and/or longer time periods, this information may not be particularly relevant for marine managers or policy makers who require accurate local scale information, particularly on magnitude, to assess likely impacts. For other research applications, such as studies of large-scale climate variability, MHW metrics may require further modification based on the resolution of datasets being used. For example, large-scale gridded data products can be used to examine the size-frequency of MHWs and their intrinsic climatic properties by setting lower thresholds to capture enough discrete warming events for

391 statistical analysis (Scannell et al. in review). While this approach is applicable for large-
392 scale MHW pattern recognition, it does not resolve the frequency of shorter and more intense
393 MHWs that would benefit from high temporal and spatial resolution data.

394 3. Monitoring and forecasting marine heatwaves

395 The three regional examples provided in section 2 demonstrate that both large and small
396 MHWs are detected in observational data based on our definitions (**Figure 3)**. In order to
397 identify the risk of MHW impacts on biological activity, the thermal thresholds of the
398 performance of different biological traits must also be known, and is the subject of ongoing
399 research. Although impacts on marine environments are still poorly understood, as detailed
400 earlier, extreme temperature events can affect species distributions and alter ecosystem
401 structure. Thus monitoring and forecasting are important and can be advanced by the use of
402 common metrics to understand and minimise potential impacts on ecological and economic
403 (e.g. fisheries) levels. Near real-time monitoring using the hierarchical classification of
404 metrics discussed here and applied to daily SST data would allow warnings to be issued when
405 areas approach or exceed their specific thresholds (Spillman 2011). For example, Coral Reef
406 Watch is based on near real-time monitoring during the warmest months of the year and is
407 used to identify areas where conditions may be approaching those conducive to coral
408 bleaching (http://coralreefwatch.noaa.gov/satellite/index.php). This early warning system can
409 inform management actions (to reduce additional stressors for example) which can be
410 implemented quickly (e.g. Beeden et al. 2012). In similar ways, this tool can be enabling as
411 an aid for fisheries managers to predict the potential impacts of increased temperature on
412 important habitats (Donnelly 2013), fish distributions (Hobday et al. 2011) and altered catch
413 rates, or whether perhaps they might be better placed to switch to different target species
414 expected to prosper under warmer conditions in the prospective areas (Mills et al. 2013) or
415 implement recovery actions when the event has concluded.
416
417 Furthermore, monitoring heatwaves can lead to a better understanding of their development,
418 characteristics and impacts. Near real-time monitoring of ocean surface temperatures based
419 on satellite data is possible, while deployment of submerged data loggers close to the
420 coastline and the use of oceanographic arrays for the open sea could provide information
421 about heat penetration depths and durations. Many of these systems are already in place, such
422 that implementing a reporting system triggered by the proposed hierarchical set of metrics

would allow characterisation of a MHW as it develops and persists, comparison to historical events, and greater insight into potential impacts.

Besides near real-time monitoring, the metrics can be used to estimate the prevalence of future MHWs. These metrics can be useful at different time scales in forecasting for the following days to weeks and for long-term projections. Using them within a forecasting framework would lead to near-term prediction of MHWs. Tools already exist for short-term and seasonal forecasting, for example Australia's Bureau of Meteorology OceanMAPS system predicts daily SSTs with a one-week lead time (www.csiro.au/bluelink/) and their POAMA model predicts monthly SSTs for the upcoming nine months (www.bom.gov.au/climate/poama2.4/poama.shtml). Including MHW metrics in the forecasting based on daily predictions would help to identify areas where MHWs may occur and actions could be implemented weeks ahead of time, including altering fisheries management boundaries (e.g. Hobday et al. 2011) and coral reef monitoring (Beeden et al. 2012).

Projections beyond the near-term could identify future MHW risk areas. Identifying risk areas would be a useful tool for Marine Protected Areas (MPAs) and spatial zoning for aquaculture. In planning MPAs, it is important not only to decide which areas are to be protected, but also where protection would be most useful. For example, protecting high diversity coral reef areas with a high probability of catastrophic disturbances in the near future, including MHWs, may be less favourable in comparison with protecting an area with less biodiversity but a low probability for disturbance (Game et al. 2008). The likelihood of an area experiencing extreme climatic events could thus be used to decide which areas should be protected and which are less resilient and prone to strong impacts with low expectations of recovery. In a similar way, decision-making processes in aquaculture zoning could include the projection of likelihood for MHWs.

4. Recommendations and conclusions

This paper has outlined the growing interest in documenting and understanding marine heatwaves. The adverse impacts of these events span a vast range of marine ecosystems. Atmospheric heatwaves have had a large research focus in recent years and a proliferation of heatwave metrics now exist, largely due to an absence of coordinated efforts in marrying the

tools and needs of physical scientists and impacts researchers. There is an opportunity for the marine community to learn from this experience, and it is on this basis that we recommend a consistent, hierarchical framework in which to measure MHWs. The three-tier framework allows for an over-arching and consistent measurement of heatwaves, while also providing flexibility in specifying additional metrics, if necessary. Regarding the use of the proposed hierarchical definition and associated metrics, we recommend the following:

1. The adoption of consistent terminology, definitions and metrics by a broad range of researchers interested in MHWs. This will facilitate comparisons between different MHW events, across seasons and at regional scales. It will also facilitate the comparison of observed events against those simulated in model projections, which will be very useful in understanding plausible future changes in MHWs.
2. The use of a flexible hierarchical system allowing for further development of descriptive indices, for particular ecosystems or species as needed by individual research goals.
3. The calculation of MHWs from the highest quality data available. Confidence in the robust detection of MHWs (and capacity to compare between events and examine spatio-temporal trends) will only be achieved with the use of high-quality datasets. Temperature data should be quality controlled, collected over adequate timescales (i.e. at least 30 years for deriving climatological baselines) and at the highest possible resolution. For example, the satellite-derived SST dataset allows for robust detection of MHWs but should be complemented with high quality *in situ* data (e.g. from coastal temperature loggers or oceanographic moorings). Daily climatological threshold time series (e.g. 90^{th} percentile) may need to be smoothed in order to extract a useful climatology from inherently variable data. Sensitivity testing on daily data suggests that a 30-day 'moving window' is appropriate for smoothing climatology from daily data.
4. To be consistent with the atmospheric heatwave literature, we recommend the 90^{th} percentile be used to define a MHW threshold and that at least five continuous days above this threshold be required to define a MHW. While 10% of days will be above this threshold, it is generally "rare" for (five) consecutive days above their relative 90^{th} percentile to occur. Shorter heat spikes may have ecological impacts in the ocean, but these are distinct features and just as a few hot air days do not make an atmospheric heatwave, a short sequence of hot ocean days (<5 days) do not represent a MHW under our definition. The use of standardised software would ensure consistency in calculating metrics, but the provision of detailed formulae (**Table 2**) may be an alternative. These

metrics can, of course, be modified to suit the specific application, but reporting of standardised metrics will greatly facilitate inter-comparison between events, locations and times.

5. Assessments of spatial and temporal variability in the occurrence of MHWs can be combined with analyses of other important aspects of the marine environment, such as biodiversity patterns (Tittensor et al. 2010), human pressures (Halpern et al. 2008), and hotspots of ocean warming (Hobday and Pecl 2014) or the velocity of climate change (Burrows et al. 2011; Sen Gupta et al. 2014). Such an approach can be used to identify regions that may be particularly susceptible to MHWs (i.e. areas subjected to intense human impacts) or regions where ecological impacts may be particularly severe (i.e. hotspots of biodiversity).

Overall, in a rapidly changing climate, the detection, characterisation, impact assessment and prediction of MHWs will become increasingly important. Marine heatwaves are an emerging area of interdisciplinary research with potential for collaborative initiatives in understanding these phenomena. A recent atmospherically driven marine heatwave in the northeast Pacific during the boreal winters of 2013-2015 had significant downstream effects on North American weather, and also disrupted northeast Pacific fisheries and coastal ecosystems (Bond et al. 2015; Hartman, 2015; Whitney 2015). This event, along with the 2003 Mediterranean Sea, 2011 Western Australia and 2012 Northwest Atlantic MHW, provide an opportunity to investigate the drivers and anomalous properties of MHWs under a hierarchical framework. We recommend that the marine scientific community adopts a coherent and consistent approach to this significant undertaking and considers how advances made in the study of atmospheric heatwaves can assist research on MHWs.

Acknowledgments

This contribution is an outcome from the working group ‘Marine Heatwaves – physical drivers and properties’ hosted at the UWA Oceans Institute by TW, DAS, NJH and ECJO. The working group received support from a University of Western Australia Research Collaboration Award, a UWA School of Plant Biology synthesis grant, and the ARC Centre of Excellence for Climate System Science (ARCCSS). This work contributes to the World Climate Research Programme (WCRP) Grand Challenge on Extremes. The workshop, and this paper, makes a contribution to the interests and activities of the International Commission on Climate of IAMAS/IUGG. This is recorded as PMEL contribution number

4403. SEP was supported by ARC grant number DE140100952, MGD by ARC grant DE150100456, LVA by ARC grant number CE110001028, DS by NERC IRF NE/K008439/1, TW by ARC grant number FT110100174. MTB was supported by NERC grant NE/J024082/1, JB acknowledges support from CE110001028, EO by ARC grant numbers FS110200029 and CE110001028, PJM by Marie Curie CIG PCIG10-GA-2011-303685 and NERC grant NE/J024082/1.

References

Alexander, L. V., Zhang, X., Peterson, T. C., Caesar, J., Gleason, B., Klein Tank, A. M. G., Haylock, M., Collins, D., Trewin, B., Rahimzadeh, F., Tagipour, A., Kumar Kolli, R., Revadekar, J. V., Griffiths, G., Vincent, L., Stephenson, D. B., Burn, J., Aguilar, E., Brunet, M., Taylor, M., New, M., Zhai, P., Rusticucci, M., Vazquez Aguirre, J. L. (2006). Global observed changes in daily climate extremes of temperature and precipitation. Journal of Geophysical Research-Atmospheres **111**: D05109, doi:10.1029/2005JD006290

Andrews, S., Bennett, S., Wernberg, T. (2014): Reproductive seasonality and early life temperature sensitivity reflect vulnerability of a seaweed undergoing range reduction. Marine Ecology Progress Series **495**: 119-129.

Beeden, R., Maynard J. A., Marshall, P. A., Heron, S. F., Willis B. L. (2012). A Framework for Responding to Coral Disease Outbreaks that Facilitates Adaptive Management. Environmental Management **49**: 1-13.

Bennett, S., Wernberg, T., Connell, S. D., Hobday, A. J., Johnson, C. R., Poloczanska E. S. (2015). The 'Great Southern Reef': Socio-ecological and economic value of Australia's neglected natural wonder. Marine and Freshwater Research. doi:10.1071/MF15232

Benthuysen J., Feng, M., Zhong, L. (2014). Spatial patterns of warming off Western Australia during the 2011 Ningaloo Niño: Quantifying impacts of remote and local forcing. Continental Shelf Research, **91**: 232-246.

Berkelmans R., De'ath G., Kininmonth S., Skirving, W. (2004). A comparison of the 1998 and 2002 coral bleaching events on the Great Barrier Reef: spatial correlation, patterns, and predictions. Coral Reefs **23**: 74-83.

Bond, N.A., Cronin, M.F., Freeland, H., Mantua, N. (2015), Causes and impacts of the 2014 warm anomaly in the NE Pacific. Geophysical Research Letters **42**: doi:10.1002/2015GL063306

Burrows, M. T., Schoeman, D. S., Buckley, L. B., Moore, P. J., Poloczanska, E. S., Brander, K. M., Brown, C. J., Bruno, J. F., Duarte, C. M., Halpern, B. S., Holding, J., Kappel, C. V., Kiessling, W., O'Conner, M. I., Pandolfi, J. M., Parmesan, C., Schwing, F. B., Sydeman, W. J, Richardson, A. J. (2011). The Pace of Shifting Climate in Marine and Terrestrial Ecosystems. Science **334**(652): DOI: 10.1126/science.1210288.

Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso, A., McPhaden, M. J., Wu, L., England, M. H., Wang, G., Guilyardi, E.,

563 Jin, F.-F. (2014). Increasing frequency of extreme El Nino events due to greenhouse
564 warming. Nature Climate Change **4**:111-116.
565 Caputi, N., Feng, M., Pearce, A., Benthuysen, J., Denham, A., Hetzel, Y., Matear, R.,
566 Jackson, G., Molony, B., Joll, L., Chandrapavan, A. (2015). Management implications
567 of climate change effect on fisheries in Western Australia. FRDC Project 2010/535.
568 Fisheries Research Report No. 260, Department of Fisheries, Western Australia, 180
569 pp.
570 Chen, K., Gawarkiewicz, G. G., Lentz, S. J., Bane, J. M. (2014), Diagnosing the warming of
571 the Northeastern U.S. Coastal Ocean in 2012: A linkage between the atmospheric jet
572 stream variability and ocean response, Journal of Geophysical Research **119**:1-10:
573 DOI:10.1002/2013JC009393
574 Cheung, W. W. L., Lam, V. W. Y., Sarmiento, J. L., Kearney, K., Watson, R., Pauly, D.
575 (2009). Projecting global marine biodiversity impacts under climate change scenarios.
576 Fish and Fisheries **10**(3): 235-251, DOI: 10.1111/j.1467-2979.2008.00315.x.
577 Coumou, D., Rahmstorf, S. (2012). A decade of weather extremes. Nature Climate Change,
578 **2**: 491-496.
579 Cowan, T., Purich, A., Perkins, S., Pezza, A., Boschat, G., Sadler, K. (2014). More frequent,
580 longer, and hotter heat waves for Australia in the Twenty-First Century. Journal of
581 Climate, **27**(15): 5851-5871
582 De'ath G., Fabricius K.E., Sweatman H., Puotinen M. (2012). The 27–year decline of coral
583 cover on the Great Barrier Reef and its causes. Proceedings of the National Academy of
584 Sciences, **109**: 17995-17999.
585 Della-Marta, P. M., Haylock, M. R., Luterbacher, J., Wanner, H. (2007), Doubled length of
586 western European summer heat waves since 1880, Journal of Geophysical Research,
587 **112**: D15103, doi:10.1029/2007JD008510.
588 Donnelly, R. J. (2013). Stewardship action plan 2013: Mitigating ecological risk in a
589 changing climate. Pro-Vision Reef Inc., Cairns Australia. Available at
590 www.provisionreed.org.au
591 Donner, S. D., Skriving, W. J., Little, C. M., Oppenheimer, M., Hoegh-Guldberg, O. (2005).
592 Global assessment of coral bleaching and required rates of adaptation under climate
593 change. Global Change Biology **11**: 2551-2265.
594 Donner, S. D. (2011). An evaluation of the effect of recent temperature variability on the
595 prediction of coral bleaching events. Ecological Applications **21**: 1718–1730.

596 Eakin, C.M., Morgan, J.A., Heron, S.F., Smith, T.B., Liu, G., Alvarez-Filip, L., Baca, B.,
597 Bartels, E., Bastidas, C., Bouchon, C., Brandt, M., Bruckner, A.W., Bunkley-Williams,
598 L., Cameron, A., Causey, B.D., Chiappone, M., Christensen, T.R.L., Crabbe, M.J.C.,
599 Day, O., de la Guardia, E., Díaz-Pulido, G., DiResta, D., Gil-Agudelo, D.L., Gilliam,
600 D.S., Ginsburg, R.N., Gore, S., Guzmán, H.M., Hendee, J.C., Hernández-Delgado,
601 E.A., Husain, E., Jeffrey, C.F.G., Jones, R.J., Jordán-Dahlgren, E., Kaufman, L.S.,
602 Kline, D.I., Kramer, P.A., Lang, J.C., Lirman, D., Mallela, J., Manfrino, C., Maréchal,
603 J.-P., Marks, K., Mihaly, J., Miller, W.J., Mueller, E.M., Muller, E.M., Orozco Toro,
604 C.A., Oxenford, H.A., Ponce-Taylor, D., Quinn, N., Ritchie, K.B., Rodríguez, S.,
605 Ramírez, A.R., Romano, S., Samhouri J.F., Sánchez, J.A., Schmahl, G.P., Shank, B.V.,
606 Skirving, W.J., Steiner, S.C.C., Villamizar, E., Walsh, S.M., Walter, C., Weil, E.,
607 Williams, E.H., Roberson,, K.W., Yusuf, Y. (2010). Caribbean Corals in Crisis: Record
608 Thermal Stress, Bleaching, and Mortality in 2005. PLoS ONE, **5**: e13969.
609 Engler, R., Randin, C.F., Thuiller, W., Dullinger, S., Zimmermann, N.E., Araujo, M.B.,
610 Pearman, P.B., Le Lay, G., Piedallu, C., Albert, C.H., Choler, P., Coldea, G., De Lamo,
611 X., Dirnbock, T., Gegout, J.-C., Gomez-Garcia, D., Grytnes, J.-A., Heegaard, E.,
612 Hoistad, F., Nogues-Bravo, D., Normand, S., Puscas, M., Sebastia, M.-T., Stanisci, A.,
613 Theurillat, J.-P., Trivedi, M.R., Vittoz, P., Guisan, A. (2011). 21st century climate
614 change threatens mountain flora unequally across Europe. Global Change Biology **17**:
615 2330-2341.
616 Fanger, P. O., (1970) Thermal comfort. Analysis and applications for environmental
617 engineering. Danish Technical Press, Copenhagen. 244pp.
618 Feng, M., McPhaden, M.J., Xie, S.-P., Hafner, J. (2013). La Niña forces unprecedented
619 Leeuwin Current warming in 2011. Scientific Reports **3**: 1277.
620 DOI:10.1038/srep01277.
621 Firth, L.B., Knights, A.M., Bell, S.S. (2011). Air temperature and winter mortality:
622 Implications for the persistence of the invasive mussel, *Perna viridis* in the intertidal
623 zone of the south-eastern United States. Journal of Experimental Marine Biology and
624 Ecology **400**: 250-256.
625 Fischer, E. M., Schar, S. (2010) Consistent geographical patterns of changes in high-impact
626 European heatwaves. Nature Geoscience **3**:398-403. DOI: 10.1038/NGEO866.
627 Frich, P., Alexander, L. V., Della-Marta, P., Gleason, B., Haylock, M., Klein Tank, A. M. G.,
628 Peterson, T. (2002). Observed coherent changes in climatic extremes during the second
629 half of the twentieth century. Climate Research **19**: 193-212.

630 Game, E. T., Watts, M. E., Woolbridge, S., Possingham, H. P. (2008): Planning for
631 persistence in marine reserves: a question of catastrophic importance. Ecological
632 Applications **18**(3): 670-680.
633 Garrabou J., Coma, R., Bensoussan, N., Bally, M., Chevaldonne, P., Cigliano, M., Diaz, D.,
634 Harmelin,, J.G., Gambi, M.C., Kersting, D.K., Ledoux, J.B., Lejeusne, C., Linares, C.,
635 Marschal, C., Perez, T., Ribes, M., Romano, J.C., Serrano, E., Teixido, N., Torrents, O.,
636 Zabala,, M., Zuberer, F., Cerrano, C. (2009). Mass mortality in Northwestern
637 Mediterranean rocky benthic communities: effects of the 2003 heat wave. Global
638 Change Biology **15**: 1090-1103.
639 Gillanders, B.M., Kingsford, M.J. (2002). Impact of changes in flow of freshwater on
640 estuarine and open coastal habitats and the associated organisms. Oceanography and
641 Marine Biology Annual Review, **40**: 233-309.
642 Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Agrosa, C., Bruno,
643 J. F, Casey, K. S., Ebert, C., Fox, H. E., Fujita, R., Heinemann, D., Lenihan, H. S.,
644 Madin, E. M. P., Perry, M. T., Selig, E. R., Spaulding, M., Steneck R. S., Watson, R.
645 (2008). A Global Map of Human Impact on Marine Ecosystems. Science **319**: 948-952.
646 Hartmann, D.L. (2015), Pacific sea surface temperature and the winter of 2014. Geophysical
647 Research Letters **42**: 1894-1902. doi:10.1002/2015GL063083
648 Helmuth, B., Mieszkowska, N., Moore, P.J., Hawkins, S.J. (2006). Living on the edge of two
649 changing worlds: Forecasting the responses of intertidal ecosystems to climate change.
650 Annual Review of Ecology, Evolution, and Systematics **37**: 373-404.
651 Hobday, A. J., Pecl, G. T. (2014). Identification of global marine hotspots: sentinels for
652 change and vanguards for adaptation action. Reviews in Fish Biology and Fisheries **24**:
653 415-425. DOI 10.1007/s11160-013-9326-6.
654 Hobday, A. J., Hartog, J., Spillman, C., Alves, O. (2011). Seasonal forecasting of tuna habitat
655 for dynamic spatial management. Canadian Journal of Fisheries and Aquatic Sciences
656 **68**: 898–911.
657 Hodgkinson, J. A., Hobday A. J., Pinkard, E. A. (2014). Climate adaptation in Australia's
658 resource-extraction industries: ready or not? Regional Environmental Change **14**(4):
659 1663-1678 DOI 10.1007/s10113-014-0618-8.
660 IPCC (2012). *Managing the Risks of Extreme Events and Disasters to Advance Climate*
661 *Change Adaptation. A Special Report of Working Groups I and II of the*
662 *Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge,
663 UK, and New York, NY, USA.

664 IPCC (2013). Summary for Policymakers. In: Climate Change 2013: The Physical Science
665 Basis. Contribution of Working Group I to the Fifth Assessment Report of the
666 Intergovernmental Panel on Climate Change [Stocker, T. F., D. Qin, G.-K. Plattner, M.
667 Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P. M. Midgley (eds.)].
668 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA
669 Jentsch, A., Kreyling, J., Beierkuhnlein, C. (2007). A new generation of climate-change
670 experiments: events, not trends. Frontiers in Ecology and Environment **5**:365-374.
671 Lotze, H. K., Worm, B., Sommer, U. (2001): Strong bottom-up and top-down control of early
672 life stages of macroalgae. Limnology and Oceanography **46**: 749–757.
673 Marba, N., Duarte, C.M. (2010). Mediterranean warming triggers seagrass (*Posidonia*
674 *oceanica*) shoot mortality. Global Change Biology **16**: 2366-2375.
675 Marbà, N., Jorda, G., Agusti, S., Girard C., Duarte C. M. (2015). Footprints of climate
676 change on Mediterranean Sea biota. Frontiers in Marine Science
677 DOI=10.3389/fmars.2015.00056.
678 Mayer, M., Hoppe, P. (1987). Thermal comfort of man in different urban environments.
679 Theoretical and Applied Climatology **38,** 43-49.
680 Maynard J.A., Turner, P.J., Anthony, K.R.N., Baird, A.H., Berkelmans, R., Eakin, C.M.,
681 Johnson, J., Marshall, P.A., Packer, G.R., Rea, A., Willis, B.L. (2008). ReefTemp: An
682 interactive monitoring system for coral bleaching using high-resolution SST and
683 improved stress predictors. Geophysical Research Letters, 35, L05603.
684 Meehl, G.A., Tebaldi, C. (2004) More Intense, More Frequent, and Longer Lasting Heat
685 Waves in the 21st Century. Science, **305**: 994-997.
686 Mills, K.E., Pershing, A.J., Brown, C.J., Yong, C., Fu-Sung, C., Holland, D.S., Lehuta, S.,
687 Nye, J.A., Sun, J.C., Thomas, A.C., Wahle, R.A. (2013). Fisheries Management in a
688 Changing Climate Lessons from the 2012 Ocean Heat Wave in the Northwest Atlantic.
689 Oceanography **26**: 191-195.
690 Palumbi, S. R., Barshis, D. J., Traylor-Knowles, N., Bay, R. A. (2014) Mechanisms of reef
691 coral resistance to future climate change. Science **344**: 895-898.
692 Parmesan, C. , Yohe, G. (2003). A globally coherent fingerprint of climate change impacts
693 across natural systems. Nature, **421**: 37-42.
694 Pearce, A., Lenanton, R., Jackson, G., Moore, J., Feng, M., Gaughan, D. (2011). The "marine
695 heat wave" off Western Australia during the summer of 2010/11. Fisheries Research
696 Report No. 222. Department of Fisheries, Western Australia. 40pp.

697 Perkins, S.E. (2011). Biases and model agreement in the projections of climate extremes over
698 the tropical Pacific. Earth Interactions, **15**: 1-36. DOI: 10.1175/2011EI395.1
699 Perkins, S. E., Alexander, L. V. (2013). On the measurement of heat waves. Journal of
700 Climate **26**: 4500–4517. doi: http://dx.doi.org/10.1175/JCLI-D-12-00383.1
701 Perkins, S. E., Alexander, L. V., Nairn, J. (2012). Increasing frequency, intensity and
702 duration of observed global heat waves and warm spells. Geophysical Research Letters,
703 **39**: 20, doi:10.1029/2012GL053361
704 Poloczanska, E. S., C. J. Brown, W. J. Sydeman, W. Kiessling, D. S. Schoeman, P. J. Moore,
705 K. Brander, J. F. Bruno, L. B. Buckley, M. T. Burrows, C. M. Duarte, B. S. Halpern, J.
706 Holding, C. V. Kappel, M. I. O'Connor, J. M. Pandolfi, C. Parmesan, F. Schwing, S. A.
707 Thompson and A. J. Richardson (2013). Global imprint of climate change on marine
708 life. Nature Climate Change: DOI: 10.1038/NCLIMATE1958.
709 Reynolds, R.W., Rayner, N.A., Smith, T.M., Stokes D.C., Wang W. (2002) An improved in
710 situ and satellite SST analysis for climate. Journal of Climate, **15**: 1609-1625.
711 Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., Schlax, M. G. (2007)
712 Daily High-Resolution-Blended Analyses for Sea Surface Temperature. Journal of
713 Climate, **20**: 5473-5496.
714 Richardson, A.J., Poloczanska, E.S. (2008). Under-resourced, under threat. Science, **320**:
715 1294-1295.
716 Rosenzweig, C., Karoly, D., Vicarelli, M., Neofotis, P., Wu, Q., Casassa, G., Menzel, A.,
717 Root, T.L., Estrella, N., Seguin, B., Tryjanowski, P., Liu, C., Rawlins, S., Imeson, A.
718 (2008). Attributing physical and biological impacts to anthropogenic climate change.
719 Nature, **453**: 353-357.
720 Russo, S., Dosio, A., Graversen, R. G., Sillmann, J., Carrao, H., Dunbar, M. B., Singleton,
721 A., Montagna, P., Barbola, P., Vogt, J. V. (2014), Magnitude of extreme heat waves in
722 present climate and their projection in a warming world. Journal of Geophysical
723 Research Atmosphere, **119**: 12,500–12,512, doi:10.1002/2014JD022098.
724 Santelices, B. (1990) Patterns of reproduction, dispersal and recruitment in seaweeds.
725 Oceanography and Marine Biology Annual Review **28**: 177–276.
726 Scannell, H.A., Pershing, A. J., Alexander, M. A., Thomas, A. C., Mills, K. E. (in review).
727 Frequency of marine heatwaves in the North Atlantic and North Pacific since 1950.
728 Schoetter, R., Cattiaux J., Douville, H. (2014). Changes of western European heat wave
729 characteristics projected by the CMIP5 ensemble. Climate Dynamics, DOI
730 10.1007/s00382-014-2434-8

731 Selig, E.R., Casey, K.S., Bruno, J.F. (2010). New insights into global patterns of ocean
732 temperature anomalies: implications for coral reef health and management. Global
733 Ecology and Biogeography **19**: 397-411.
734 Sen Gupta, A., Brown, J. N., Jourdain, N. C., van Sebille, E., Ganachaud, A., Vergés A.
735 (2015). Episodic and non-uniform shifts of thermal habitats in a warming ocean, Deep
736 Sea Research Part II: Topical Studies in Oceanography, **113**: 59–72,
737 doi:10.1016/j.dsr2.2013.12.002.
738 Smale, D. A., Wernberg, T. (2009). Satellite-derived SST data as a proxy for water
739 temperature in nearshore benthic ecology. Marine Ecology Progress Series **387**:27-37
740 Smale, D. A., Wernberg, T. (2013). Extreme climatic event drives range contraction of a
741 habitat-forming species. Proceedings of the Royal Society of London, Series B:
742 Biological Sciences **280**:20122829): http://dx.doi.org/10.1098/rspb.2012.2829.
743 Smith, M.D. (2011). The ecological role of climate extremes: current understanding and
744 future prospects. Journal of Ecology **99**: 651-655.
745 Sorte, C.J.B., Fuller, A., Bracken, M.E.S. (2010). Impacts of a simulated heat wave on
746 composition of a marine community. Oikos **119**: 1909-1918.
747 Spillman, C., Alves, O. (2009). Dynamical seasonal prediction of summer sea surface
748 temperatures in the Great Barrier Reef. Coral Reefs **28**: 197-206.
749 Spillman, C. (2011). Operational real-time seasonal forecasts for coral reef management.
750 Journal of Operational Oceanography **4**(1): 13-22.
751 Steadman, R. G. (1984). A universal scale of apparent temperature. Journal of Climate and
752 Applied Meteorology, **23,** 1674–1687
753 Tittensor, D. P., Mora, C., Jetz, W., Lotze, H. K., Ricard, D., Vanden Berghe, E., Worm, B.
754 (2010). Global patterns and predictors of marine biodiversity across taxa. Nature **466**:
755 1098–1101 doi:10.1038/nature09329.
756 Trenberth, K. (2012). Framing the way to relate climate extremes to climate change. Climatic
757 Change **115**:283-290.
758 Wernberg, T., Smale, D.A., Tuya, F., Thomsen, M.S., Langlois, T.J., de Bettignies, T.,
759 Bennett S., Rousseaux, C.S. (2013). An extreme climatic event alters marine ecosystem
760 structure in a global biodiversity hotspot. Nature Climate Change, **3**: 78-82.
761 Whitney, F.A. (2015), Anomalous winter winds decrease 2014 transition zone productivity in
762 the NE Pacific, Geophysical Research Letters, **42**: 428-431,
763 doi:10.1002/2014GL062634

764 Zhang, X., Hegerl, G., Zwiers, F.W., Kenyon, J. (2005). Avoiding inhomogeneity in
765 percentile-based indices of temperature extremes. Journal of Climate **18**(11): 1641-
766 1651
767 Zhang, X., Alexander, L., Hegerl, G. C., Jones, P., Klein Tank, A., Peterson, T. C., Trewin,
768 B., Zwiers, F. W. (2011). Indices for monitoring changes in extremes based on daily
769 temperature and precipitation data. WIREs Climate Change doi:10.1002/wcc.147.
770 Zinke, J., Rountrey, A., Feng, M., Xie, S.-P., Dissard, D., Rankenburg, K., Lough, J. M.,
771 McCulloch, M. T. (2014). Corals record long-term Leeuwin current variability
772 including Ningaloo Nino/Nina since 1795. Nature Communications **5**:3607.
773

774 **Table 1:** Examples of metrics commonly used to describe warming events in ecological
775 studies.

Metric	Description	Example
Maximum Temperature (°C)	Maximum temperature observed during a heatwave event. E.g. 30°C.	Berkelmans et al. 2004
Temperature Anomaly (°C)	Deviation from long-term mean (most often monthly mean). E.g. 3.5° C above average.	Sorte et al. 2010; Wernberg et al. 2013; Smale and Wernberg 2013
Thermal Stress Anomaly (e.g. weeks)	Temperature deviation above a threshold value (rather than the mean value), summed over some period of time (e.g. weeks). E.g. TSA = 45°C over 10 weeks.	Selig et al. 2010
Degree Heating Weeks (°C-weeks)	Degree Heating Weeks (DHW) reflect the accumulation of heat stress by integrating SST anomalies in excess of a threshold over a period of weeks. In corals, thermal stress occurs when sea surface temperatures exceed a certain threshold (usually defined as ~1°C above the maximum climatological mean), and so DHWs are calculate as the sum of SST anomalies above the 1°C threshold over a number of weeks (e.g. 12 weeks).	Eakin et al. 2010 Donner 2011
Degree Heating Days (°C days)	The degree heating days (DHD) value is the summed positive deviations of daily mean sea surface temperatures (x(t)) from the climatology of long-term mean summer temperatures (LMST), for a specified period (e.g. summer, December 1st to February 28th in the Southern Hemisphere). $$DHD = \sum (x(t) - LMST)$$	Maynard et al. 2008
Heating rate (°C/day)	Heating rate (HR) is defined as $\frac{DHD}{ND}$ where DHD is degree heating days as defined above, and ND is the number of days in which daily mean sea surface temperatures (x(t)) have exceeded the long-term mean summer temperatures. That is, HR is the mean rate at which DHD have accumulated throughout a period of time (e.g. summer, December 1st to February 28th in the Southern Hemisphere).	Maynard et al. 2008

776
777

Table 2. Hierarchical classification of metrics to characterise marine heat waves (MHW). All definitions assume that daily SST data, T, and that a MHW has a discrete start day and end day. Note that we write T both as a function of time t, $T(t)$, and as a function of year y and day-of-year d, $T(y,d)$.

	Name	Definition		Units
	Climatology	T_m: The climatological mean, calculated over a reference period, to which all values are relative	$T_m(j) = \sum_{y=y_s}^{y_e} \sum_{d=j-5}^{j+5} \frac{T(y,d)}{11(y_e - y_s + 1)}$ where j is day of year, y_s and y_e are the start and end of the climatological base period respectively, and T is the daily SST on day d of year y	°C
	Threshold*	$T_{\%}$: The seasonally varying temperature value that defines a MHW (e.g. T_{90} is the 90th percentile value based on the baseline periods)	$T_{90}(j) = P_{90}(X)$ where P_{90} is the 90^{th} percentile and $P_{90}(X)$ where $X = \{T(y,d) \mid y_s <= y <= y_e, j-5 <= d <= j+5\}$	°C
	Start and end of MHW	t_e, t_s: dates on which a MHW begins and ends.	t_s is the time, t, where $T(t) > T_{90}(j)$ and $T(t-1) < T_{90}(j)$. t_e is the time, t, where $t_e > t_s$ and $T(t) < T_{90}(j)$ and $T(t-1) > T_{90}(j)$ For MHWs, $t_e - t_s \geq 5$, and where gap ≤2 days (see text)	days
Primary	Duration	D: Consecutive period of time that temperature exceeds the threshold	$D = t_e - t_s$	days
	Intensity (max/mean/variance)	i_{max}: highest temperature anomaly value during the MHW i_{mean}: mean temperature anomaly during the MHW i_{var}: variation in intensity of the MHW over the duration	$i_{max} = \max(T(t) - T_m(j))$ $i_{mean} = \overline{T(t) - T_m(j)}$ $i_{var} = \sigma_{T(t)}$ where $t_s \leq t \leq t_e$, $j(t_s) \leq j \leq j(t_e)$, σ is the standard deviation, and the overbar indicates the time mean.	°C
Secondary	Rate measures	r_{onset}: time from the onset of the MHW to the maximum intensity. $r_{decline}$: time from	$r_{onset} = \frac{i_{max} - (T(t_s-1) - T_m(j-1))}{t_{max} - (t_s - 1)}$	°C/day

		the maximum intensity to the end of the MHW.	$r_{decline} = \frac{i_{max} - \left(T(t_e) - T_m(j)\right)}{t_e - t_{max}}$ where t_{max} is the time of MHW_{imax}	
	Cumulative measure	i_{cum}: sum of daily intensity anomalies. Note that the integral omits t_e which is below the T_{90} threshold.	$i_{cum} = \int_{t_s}^{t_{e-1}} (T(t) - T_m(j))dt$	°C days
	Spatial extent	A: Area of ocean meeting the MHW definition L: Length of coastline for the MHW	A = area over which MHW detected L = length of coast where MHW detected	km^2 km
Tertiary	Preconditioning factors	Factors such as time of year relative to the onset of the MHW, or periods of above mean temperature preceding the MHW may lead to greater impacts.	n/a	Various – specific to study system

782
783
784

Table 3. Qualitative comparison of different temperature data sources and their suitability to provide primary, secondary and tertiary marine heat wave metrics for sea surface temperature (SST). Relative scores for each option are in the range 1 to 4, where 1 indicates that only low resolution metrics can be derived and 4 indicates that high resolution metrics can be derived (N/A indicates no utility). The quantities maximum MHW intensity (i_{max}) and cumulative effect (i_{cum}) are defined in **Figure 1** and **Table 2**. Preconditioning is defined as the conditions that facilitate the onset of the MHW. Continuous data generally allow an understanding of environmental conditions leading up to the event.

Data source	Metrics						Other considerations			
	Primary			Secondary		Tertiary				
	Duration (D) [days]	i_{max} [°C]	i_{mean}, i_{cum} [°C days]	Rate of event onset/decay [°C/day]	Spatial area (A) [km^2]	Preconditioning	Length of records	Temporal resolution	Quality control	Data consistency
In situ temperatures (e.g. loggers)	4	4	4	4	1 (if multiple loggers), else N/A	2	High	High	Low	Low/ Med
Satellite SST	3	3	3	4	3	3	Med	High	High	Low/Med
Argo floats (NB: gridded products do not provide SST)	N/A	N/A	1	1	2	2	Low	Low	High	Med/High
Reconstructed monthly data (e.g. ERSST, HadISST)	2	2	N/A	2	2	2	High	Low	High	Med
Palaeo-proxy SST (seasonal to annual records, e.g. coral cores)	N/A	N/A	N/A	N/A	N/A	N/A	High	Low	Low	Low/Med
Global Climate Models (e.g. daily SST fields)	2	2	2	3	2	2	High	Low	N/A	High
Re-analysis SST products (e.g. BRAN)	3	3	3	3	3	3	Med	High	Low/ Med	Med/High
Regional Ocean Models (e.g. OFAM)	3	3	3	3	2	2	Low	High	N/A	High
Traditional Ecological Knowledge, citizen science, and anecdotal information	N/A	1	N/A	N/A	N/A	1	Med/High	Low	Low	Low

Figure Legends

Figure 1. Schematic of metrics used to define a marine heatwave (MHW). (a) Threshold values for each location for each day of the year are defined based on the 90th percentile value. (b) These percentile values vary through the year (dashed line), as does the climatological mean (solid line). (c) Short duration heat spikes less than five days are not MHWs. A temperature event that is at least five days or longer than this minimum duration is defined according to duration (MHW_D) above the threshold value, intensity (i_{max}, temperature above the climatological mean) and the rate of temperature increase (r_{onset}) and decrease ($r_{decline}$) during the event. The mean event intensity (open circle, i_{mean}) is the mean intensity during the MHW, while i_{cum} (shading) is the sum of daily intensities during the MHW. The start and end days of the MHW are represented by t_s and t_e respectively.

Figure 2. Taxonomy of heat waves as distinguished by the metrics duration (D), maximum intensity (i_{max}), cumulative intensity (i_{cum}), and rate of onset (r_{onset}). A marine heat wave (MHW) with regular warming onset and decline (panel **a**) can be distinguished from one with similar duration and maximum intensity but asymmetric warming (panel **b**) by the cumulative intensity metric (i_{cum}). This asymmetric MHW (b) is distinguished from one with a slow onset and rapid decline (panel **c**) by the rate of warming (r_{onset}) metric. A lower intensity MHW (panel **d**) is distinguished by its maximum intensity (i_{max}), while a short MHW (panel **e**) is distinguished by its duration (*D*). The dashed line indicates the threshold value. Arrows between the plots indicate the major change (Δ) between the plots. Index values are indicative only in this schematic.

Figure 3. First row: Sea surface temperatures (SST) anomaly on the peak day of three marine heatwaves (MHW) discussed in the text. (a) Western Australia 2011, (b) northern Mediterranean 2003, (c) northwest Atlantic 2012. Dots show the locations from which 1/4° resolution time series of SST were extracted from NOAA OI SST for the detection of MHWs in each case study region. Second row: The SST climatology (blue), 90th percentile MHW threshold (green), and SST time series (black) for each MHW at each location. The red filled area indicates the period of time associated with the identified MHW, while shaded orange indicates other MHWs identified over the year. Third row: The duration (D) of each MHW detected in the time series from each location, with every tenth event identified on the upper

x-axis. Fourth row: As for the third row, but illustrating maximum intensity (i_{max}) of each MHW event in each location. Fifth row: As for the third row, but illustrating cumulative intensity (i_{cum}) of each MHW event from each location. The WA and northwest Atlantic MHWs are the largest by maximum intensity, such that the red and yellow bars are the same. The northwest Atlantic event is not the largest according to duration or cumulative intensity, but the red bar obscures the yellow bar since they are so close in time.

Figure 4. Temperature time series during the twelve months bounding the 2011 'Ningaloo Niño' marine heat wave (shaded area: December 2011 to April 2012) off the coast of Western Australia as measured by four different data sources; weekly and monthly Reynolds SST (29.5-30.5°S; 114.5-115.5°E), daily satellite SST (29.5-30.5°S; 114.5-115.5°E), and an hourly *in situ* logger from Jurien Bay (30 18.5 °S 114 58.3 °E).




Figure 1. Schematic of metrics used to define a marine heatwave (MHW). (a) Threshold values for each location for each day of the year are defined based on the 90th percentile value. (b) These percentile values vary through the year (dashed line), as does the climatological mean (solid line). (c) Short duration heat spikes less than five days are not MHWs. A temperature event that is at least five days or longer than this minimum duration is defined according to duration (MHW_D) above the threshold value, intensity (i_{max}, temperature above the climatological mean) and the rate of temperature increase (r_{onset}) and decrease ($r_{decline}$) during the event. The mean event intensity (open circle, i_{mean}) is the mean intensity during the MHW, while i_{cum} (shading) is the sum of daily intensities during the MHW. The start and end days of the MHW are represented by t_s and t_e respectively.




Figure 2. Taxonomy of heat waves as distinguished by the metrics duration (D), maximum intensity (i_{max}), cumulative intensity (i_{cum}), and rate of onset (r_{onset}). A marine heat wave (MHW) with regular warming onset and decline (panel **a**) can be distinguished from one with similar duration and maximum intensity but asymmetric warming (panel **b**) by the cumulative intensity metric (i_{cum}). This asymmetric MHW (b) is distinguished from one with a slow onset and rapid decline (panel **c**) by the rate of warming (r_{onset}) metric. A lower intensity MHW (panel **d**) is distinguished by its maximum intensity (i_{max}), while a short MHW (panel **e**) is distinguished by its duration (*D*). The dashed line indicates the threshold value. Index values are indicative only in this schematic.




Figure 3. First row: Sea surface temperatures (SST) anomaly on the peak day of three marine heatwaves (MHW) discussed in the text. (a) Western Australia 2011, (b) northern Mediterranean 2003, (c) northwest Atlantic 2012. Dots show the locations from which 1/4° resolution time series of SST were extracted from NOAA OI SST for the detection of MHWs in each case study region. Second row: The SST climatology (blue), 90th percentile MHW threshold (green), and SST time series (black) for each MHW at each location. The red filled area indicates the period of time associated with the identified MHW, while shaded orange indicates other MHWs identified over the year. Third row: The duration (D) of each MHW detected in the time series from each location, with every tenth event identified on the upper x-axis. Fourth row: As for the third row, but illustrating maximum intensity (i_{max}) of each MHW event in each location. Fifth row: As for the third row, but illustrating cumulative intensity (i_{cum}) of each MHW event from each location. The WA and northwest Atlantic MHWs are the largest by maximum intensity, such that the red and yellow bars are the same. The northwest Atlantic event is not the largest according to duration or cumulative intensity, but the red bar obscures the yellow bar since they are so close in time.




Figure 4. Temperature time series during the twelve months bounding the 2011 'Ningaloo Niño' marine heat wave (shaded area: December 2011 to April 2012) as measured by four different data sources; weekly and monthly Reynolds SST (29.5-30.5°S; 114.5-115.5°E), daily satellite SST (29.5-30.5°S; 114.5-115.5°E), and an hourly *in situ* logger from Jurien Bay (30 18.5 °S 114 58.3 °E).